

Supplementary Material for

**Elemental Analysis of Chamber Organic Aerosol
Using an Aerodyne High-Resolution Aerosol Mass Spectrometer**

P. S. Chhabra¹, R. C. Flagan^{1,2}, and J. H. Seinfeld^{1,2}

¹Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA

²Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA

Correspondence to: J. H. Seinfeld
(seinfeld@caltech.edu)

Table S1: Experimental conditions and elemental ratios for each experiment. Elemental ratios given are those at the time of maximum O/C.

Table S2: Calibration factors and estimated uncertainties for O/C, H/C, N/C and OM/OC as determined by Aiken et al. (2008).

Table S3: Ratios of particle-phase signals of CO^+ to CO_2^+ . The particle phase signals of H_2O^+ , OH^+ and O^+ were taken to be 22.5%, 5.625%, and 0.90% of the particle phase CO_2^+ signal.

Table S4: O/C and H/C ratios of α -pinene ozonolysis SOA determined from offline analysis by Yu et al. (1999, Table XI).

Figure S1: $\text{O}/\text{C}_{\text{HR}}$ and O/C_{44} for glyoxal uptake SOA.

Figure S2: High-resolution spectra of N-containing ions for glyoxal uptake SOA.

Figure S3: $\text{O}/\text{C}_{\text{HR}}$ and O/C_{44} for α -pinene ozonolysis SOA.

Figure S4: $\text{O}/\text{C}_{\text{HR}}$ and O/C_{44} for isoprene SOA formed under low- NO_x conditions.

Figure S5: High-resolution spectrum of m/z 91 for isoprene high- NO_x SOA.

Figure S6: $\text{O}/\text{C}_{\text{HR}}$ and O/C_{44} for isoprene SOA formed under high- NO_x conditions.

Figure S7: $\text{O}/\text{C}_{\text{HR}}$ and O/C_{44} for single-ring aromatic SOA.

Figure S8: $\text{O}/\text{C}_{\text{HR}}$ and O/C_{44} for naphthalene SOA formed under low- NO_x conditions.

Figure S9: $\text{O}/\text{C}_{\text{HR}}$ and O/C_{44} for naphthalene SOA formed under high- NO_x conditions.

High-resolution AMS spectra will be available online at <http://cires.colorado.edu/jimenez-group/HRAMSsd/>.

Expt. #	VOC System	Experiment Type	ΔM_o (Max) ($\mu\text{g}/\text{m}^3$)	O/C (Max)	H/C	N/C	OM/OC
1	Glyoxal uptake	humid	31.80	1.12	1.55	0.01	2.64
2	Glyoxal uptake	humid	68.30	1.12	1.54	0.01	2.63
3	Glyoxal uptake	humid	NA	1.15	1.52	0.02	2.68
4	α -pinene + O ₃	no H ₂ O ₂ ; dry	59.70	0.38	1.48	0.00	1.63
5	α -pinene + O ₃	no H ₂ O ₂ ; humid	87.50	0.48	1.46	0.01	1.76
6	α -pinene + O ₃	H ₂ O ₂ ; dry	127.10	0.40	1.48	0.00	1.68
7	α -pinene + O ₃	H ₂ O ₂ ; humid	192.46	0.45	1.44	0.01	1.72
8	Isoprene + OH	low NO _x	3.71	0.58	1.66	0.00	1.92
9	Isoprene + OH	low NO _x	7.00	0.58	1.69	0.00	1.92
10	Isoprene + OH	low NO _x	10.47	0.60	1.57	0.00	1.94
11	Isoprene + OH	high NO _x	1.35	0.60	1.40	0.03	1.95
12	Isoprene + OH	high NO _x	4.27	0.62	1.49	0.04	2.00
13	Isoprene + OH	high NO _x	11.83	0.65	1.49	0.04	2.04
14	Toluene + OH	low NO _x	141.45	0.74	1.39	0.00	2.10
15	Toluene + OH	high NO _x	50.26	0.72	1.38	0.07	2.15
16	<i>m</i> -xylene + OH	low NO _x	190.40	0.60	1.54	0.00	1.93
17	<i>m</i> -xylene + OH	high NO _x	52.04	0.66	1.48	0.08	2.09
18	Naphthalene + OH	low NO _x	12.15	0.62	0.95	0.00	1.92
19	Naphthalene + OH	low NO _x	44.91	0.73	0.93	0.00	2.05
20	Naphthalene + OH	low NO _x , nucleation	47.17	0.62	0.85	0.00	1.91
21	Naphthalene + OH	low NO _x	53.12	0.73	0.86	0.00	2.06
22	Naphthalene + OH	low NO _x	201.75	0.60	0.82	0.00	1.87
23	Naphthalene + OH	high NO _x	5.90	0.65	1.01	0.07	2.03
24	Naphthalene + OH	high NO _x	39.02	0.60	0.88	0.04	1.92
25	Naphthalene + OH	high NO _x , nucleation	39.26	0.51	0.90	0.04	1.79
26	Naphthalene + OH	high NO _x	75.43	0.54	0.80	0.02	1.82

Table S1

Ratio	Calibration Factor	Measurement Uncertainty
O/C	0.75	31%
H/C	0.91	10%
N/C	0.96	22%
OM/OC	-	6%

Table S2

System	Organic CO ⁺ /CO ₂ ⁺ Estimate
α-pinene-O ₃	0.98
Glyoxal Uptake	5.00
Isoprene-OH	1.00
Aromatic-OH	1.03
Naphthalene-OH	1.17

Table S3

Product Name	Formula	Molar Yield		
		6-9-98a	6-9-98b	6-17-98a
Pinic Acid	C ₉ H ₁₄ O ₄	1.8	3.9	2.8
Norpinic Acid	C ₈ H ₁₂ O ₄	0.08	0.09	0.05
Hydroxy pinonaldehydes	C ₁₀ H ₁₆ O ₃	2.4	1.1	2
Pinonic Acid	C ₁₀ H ₁₆ O ₃	1.7	1.6	1.3
Norpinonic Acid and Isomers	C ₉ H ₁₄ O ₃	2.1	4.8	2.8
Pinonaldehyde	C ₁₀ H ₁₆ O ₂	0.8	0.3	0.9
Norpinonaldehyde	C ₉ H ₁₄ O ₂	0.1	0.2	0.2
Hydroxy pinonic acid	C ₁₀ H ₁₄ O ₄	2.1	1.3	2.1
A13	C ₁₀ H ₁₆ O ₃	0.08	0.12	0.1
A14	C ₁₀ H ₁₄ O ₃	0.55	0.48	0.8
O/C		0.34	0.36	0.35
H/C		1.58	1.56	1.57

Table S4




Figure S1




Figure S2







Figure S3




Figure S4




Figure S5




Figure S6




Figure S7




Figure S8




Figure S9

References

Aiken, A. C., Decarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, 42, 4478–4485, doi:10.1021/Es703009q, 2008.

Yu, J. Z., Cocker, D. R., Griffin, R. J., Flagan, R. C., and Seinfeld, J. H.: Gas-phase ozone oxidation of monoterpenes: Gaseous and particulate products, *J. Atmos. Chem.*, 34, 207–258, 1999.